



**This material is the copyright of the original publisher.  
Unauthorised copying and distribution is prohibited.**



### **Terms and Conditions for Use of PDF**

The provision of PDFs for authors' personal use is subject to the following Terms & Conditions:

The PDF provided is protected by copyright. All rights not specifically granted in these Terms & Conditions are expressly reserved. Printing and storage is for scholarly research and educational and personal use. Any copyright or other notices or disclaimers must not be removed, obscured or modified. The PDF may not be posted on an open-access website (including personal and university sites).

The PDF may be used as follows:

- to make copies of the article for your own personal use, including for your own classroom teaching use (this includes posting on a closed website for exclusive use by course students);
- to make copies and distribute copies (including through e-mail) of the article to research colleagues, for the personal use by such colleagues (but not commercially or systematically, e.g. via an e-mail list or list serve);
- to present the article at a meeting or conference and to distribute copies of such paper or article to the delegates attending the meeting;
- to include the article in full or in part in a thesis or dissertation (provided that this is not to be published commercially).

# Spontaneous Reports and Pharmacogenetics

## The Role of the Pharmacovigilance Centre

A commentary, 'Pharmacogenetics and Pharmacovigilance',<sup>[1]</sup> was recently published concerning our study<sup>[2]</sup> in the same issue of the journal. The goal of our study was to gain insight into the feasibility of selecting and genotyping patients based on adverse drug reaction (ADR) reports received by the Netherlands Pharmacovigilance Centre. The main question posed was whether pharmacovigilance centers can provide the appropriate clinical and scientific information needed for the decision to genotype a patient who has experienced an ADR that is possibly due to a genetic polymorphism. Pharmacovigilance centres assess suspected ADRs reported by health professionals and patients. A lack of certainty about the causal relationship should not prevent health professionals and patients from reporting these ADRs; a report of a suspected ADR rarely implies a certain causal relationship between the suspected drug and the reported event. In clinical practice, the etiology of a clinical event is not always known, even after thorough examination and testing. ADRs are no exception. In addition, the event may have a multi-factorial etiology. One of these factors may be the existence of an aberrant pharmacokinetic mechanism that may explain an increased likelihood of a particular patient developing ADRs.

Assessors of ADR reports sometimes have access to specific information that should preferably also be shared with those who have actual contact with the patients. This usually applies not only to the number of reports in the (national) ADR database and information from the literature, but assessors may also know of evidence of a possible mechanism that may contribute to the occurrence of an ADR. Limiting the role of a pharmacovigilance centre to merely a reporting centre, and withholding this information from a

treating physician, could be considered as unethical. A striking example of the duty to inform patients about a genetic variant that causes serious adverse effects is the case of pseudocholinesterase. In 1956, Lehman and Ryan<sup>[3]</sup> discovered a familial factor in the 'idiopathic' low pseudocholinesterase level associated with suxamethonium chloride toxicity. Today, no-one would overlook a serious ADR report involving suxamethonium chloride or mivacurium chloride without recommending pharmacogenetic testing. What about a report on azathioprine myelotoxicity<sup>[4]</sup> – should a pharmacovigilance centre suggest a thiopurine methyltransferase (TPMT) test? Would just registration – without reporting back our knowledge – be ethical? In our view, pharmacovigilance centres do indeed have an 'advisory and educational role'.

Spontaneous reporting systems primarily have a signalling function. The causal relationship between the reported ADR and the suspected drug is rarely certain. Nevertheless, it is the task of a pharmacovigilance centre to reduce this uncertainty as much as possible. One way of doing this is by asking for additional information from the reporter about the circumstances around the ADR or any additional testing that has been carried out. For all parties involved, it is important to investigate all factors that may have contributed to the occurrence of the ADR. Also, the existence of a pharmacogenetic cause should be considered. Once a polymorphism has been established, the patient should be informed.

Providing information to healthcare professionals should be distinguished from recommending genotyping. The feedback we give to healthcare professionals merely suggests that factors such as pharmacogenetic involvement might have a role to play. Having knowledge of a possible mechanism does not imply that pharmacogenetic tests have to be carried out. Clinical judgement plays a key role in this process. Taking into consideration all other possible reasons for the ADR, it is up to the physician to decide whether or not to have the patient genotyped. In addition, knowledge of an enzyme deficiency can help the physician to choose a better medication (in the correct dosage) for the patient next time.

Another point of criticism raised in the commentary<sup>[1]</sup> referred to the exclusion of patients with allergic reactions. In patients with allergic reactions, kinetic factors very rarely play a role. As in other type B reactions, dose dependency is not likely. For this reason we decided not to include reports with these types of ADRs.

We are only just beginning to understand the role of pharmacogenetics in the pathogenesis of ADRs. Our study was a pilot study to get experience with reporting back any potential pharmacogenetic involvement in the pathogenesis of ADRs. The study has the usual limitations of a practical and naturalistic study on reported ADRs: the impossibility of obtaining blood from deceased patients retrospectively; limitation to the most prevalent pharmacogenetic factors – cytochrome P450 (CYP) 2D6, CYP2C19 and CYP2C9; the voluntary nature of patient participation; and the difficult classification and broad spectrum of the reported ADRs. As with any new methodology, its actual value only becomes evident after gaining enough experience with it. Creating awareness and informing healthcare professionals and patients about the possibility of pharmacogenetics as a causal factor

is the first step in this process. This is exactly what our study aimed to convey.

*Eugène van Puijenbroek,<sup>1</sup> Jean Conemans<sup>2</sup> and Kees van Grootheest<sup>1</sup>*

- 1 Netherlands Pharmacovigilance Centre Lareb, 's-Hertogenbosch, the Netherlands
- 2 Hospital Pharmacy ZANOB, 's-Hertogenbosch, the Netherlands

### Acknowledgements

The authors have no conflicts of interest that are directly relevant to the content of this correspondence.

### References

1. Howland H. Pharmacogenetics and pharmacovigilance. *Drug Safety* 2009; 32 (3): 265-70
2. Van Puijenbroek EP, Conemans J, Van Grootheest AC. Spontaneous ADR reports as a trigger for pharmacogenetic research: a prospective observational study in the Netherlands. *Drug Saf* 2009; 32 (3): 255-64
3. Lehman H, Ryan E. The familial incidence of low pseudocholinesterase levels. *Lancet* 1956 Jul 21; 271 (6934): 124
4. Evans WE. Pharmacogenetics of thiopurine S-methyltransferase and thiopurine therapy. *Ther Drug Monit* 2004; 26: 186-91